

[illegible]

This invention is directed to therapeutic methods involving administration of antagonists of S-nitrosothiols. S-nitrosothiols are naturally occurring compounds related to nitric oxide which are involved in the regulation of blood pressure, pain perception, control of smooth muscle tone, and numerous other functions in humans.

Nitric oxide is a major intercellular signal in humans and other animals that affects many different organ systems. Important actions caused by nitric oxide include vasodilation, smooth muscle relaxation in the gastrointestinal tract, lungs, and uterus, and many neurological functions including pain perception. Many of the actions of nitric oxide are actually caused by compounds, known as S-nitrosothiols or thionitrites, that are derived from nitric oxide and specific thiols. Some of the actions of S-nitrosothiols occur at stereospecific recognition sites that are selective for specific nitrosothiols.

One pathological condition in humans that is caused by the overproduction of nitric oxide is septic shock. The profound hypotension, tissue ischemia, and organ failure of septic shock are rare sequelae to infection yet septic shock is the leading cause of death in intensive care units. The hypotension is largely the result of massive vasodilation in the patient with hypoperfusion and organ failure, including cardiac failure, exacerbating the problem. Vasodilation is initiated by bacterial endotoxins which trigger a series of host defense mechanisms including greatly increased expression of the inducible isozyme of nitric oxide synthase

092710-054204

(iNOS). Treatment of septic shock, in addition to treatment of the underlying infection, consists of administration of intravenous fluids and vasopressors, although these therapies are less effective in septic patients than non-septic patients. Decreasing the production of nitric oxide (NO) with inhibitors of nitric oxide synthase such as N^ω-nitro-L-arginine methyl ester (L-NAME) or L-N^G-methylarginine (L-NMMA) is a logical therapy, but results in both animals and humans have not been encouraging. Heart failure and continued high mortality commonly follow such treatment.

Current treatment options for hypotension or shock from such conditions such as septic shock, toxic shock syndrome, spinal cord injury, effects of anesthetics, and anaphylaxis, etc. are limited to vasoconstricting agents that have many deleterious side effects that limit their effective therapeutic usage.

It therefore can be seen that there is a continuing need for the development of effective pharmacological treatments to counteract hypotension and shock. In particular, there is no effective treatment presently available that specifically counteracts the hypotension caused by the relative overproduction of nitric oxide and nitrosothiols as present in such conditions as septic shock. This invention has as its primary objective the fulfillment of this need.

Other objectives of the present invention include both processes, compositions and treatments useful to treat the undesired effects of nitric oxide and nitrosothiol production. Conditions other than hypotension that might benefit from inhibition of the actions of nitric oxide and nitrosothiols include, but are not limited to, chronic and acute pain, hypoxemia secondary in part to the loss of hypoxic pulmonary vasoconstriction, uterine atony, and decreased tone and peristalsis of gastrointestinal smooth muscle.

SUMMARY OF THE INVENTION

The inventors have discovered that the use of certain S-alkylthiols, including but not limited to S-ethylcysteine, S-methylcysteine, S-methylcysteamine, S-ethylcysteamine, S-ethylglutathione, S-methylglutathione, S-methylglutathione, S-methylcoenzyme A, and S-ethylcoenzyme A in treatment effective amounts as antagonists to S-nitrosothiols will significantly reduce risk of hypotension, tissue ischemia, central nervous system injury, or other acute tissue injury without significant deleterious side effects. Additional potential uses include blockage of pain conduction in peripheral nerves, spinal cord, or brain and inhibition of smooth muscle relaxation in the gastrointestinal tract, lungs, and uterus.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1 through 6 show the effects of treatments described in the examples in terms of comparison of an example of the invention antagonists to S-nitrosothiols, S-methyl-L-cysteine, to an inhibitor of nitric oxide synthase, L-NAME, in the treatment of an experimental model of septic shock.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

Preferred compounds of the present invention can generally be described as S-alkylthiols. Most preferred are those that are S-alkylthiols of certain amino acids. Particularly preferred are S-ethyl-L-cysteine, S-methyl-L-cysteine, S-ethylglutathione, S-methylglutathione, S-methylcysteamine, S-ethylcysteamine, S-methylcoenzyme A and S-ethylcoenzyme A.

The treatment of patients may be with the compounds of the present invention or their salts. Such salts include

overproduction of NO during septic shock also leads to increased production or release of S-nitrosocysteine then a specific inhibitor of S-nitrosocysteine binding might attenuate the hypotension of septic shock without affecting the potentially beneficial antimicrobial effects of nitric oxide and peroxynitrite.

We believe that a molecule structurally similar to S-nitroso-L-cysteine might be a competitor to S-nitroso-L-cysteine for the binding site. S-methyl-L-cysteine is such a compound, identical in structure to S-nitroso-L-cysteine except for the substitution of a methyl group in place of the nitroso group. In these examples, S-methyl-L-cysteine was found to inhibit the hypotensive effects of infused S-nitrosocysteine. In the present study S-methyl-L-cysteine not only attenuated the hypotension induced by LPS but did so without further impairing coronary blood flow, cardiac output, and heart rate. Total peripheral resistance returned to normal values with S-methyl-L-cysteine while treatment with L-NAME increased total peripheral resistance far above normal. The difference in the hemodynamic response between treatment with S-methyl-L-cysteine and treatment with L-NAME suggests that much of the hypotension in septic shock may be caused by overproduction of S-nitrosocysteine or some very similar compound. Inhibition of nitric oxide synthesis blocks NO-dependent vasodilation indiscriminately, leading to ischemia and organ failure even in the presence of normal mean arterial pressures. Inhibition of S-nitrosocysteine binding by S-methyl-L-cysteine selectively blocks the vasodilation caused by S-nitrosocysteine without affecting other NO-dependent vasodilations or other NO-dependent functions. Consequently, S-methyl-L-cysteine may be very useful in the treatment of septic shock.

The following examples offered to further illustrate, but not limit, the process and compositions of the present invention.

EXAMPLES

Male 16 week old Sprague-Dawley rats were anesthetized for surgical placement of intraaortic pressure transducers and Dopplar flow probes around the abdominal aorta and the left main coronary artery. After 48 hours of recovery baseline measurements of arterial blood pressure, cardiac output (as Dopplar flow), and coronary blood flow (Dopplar) were obtained in conscious unrestrained animals. The rats were then treated with intraperitoneal injections of purified lipopolysaccharide 0.5 mg/kg. After 72 hours when the induction of iNOS had produced a fall in blood pressure, measurements were repeated. Each rat then received either 25 μ mole/kg of the nitric oxide synthase inhibitor L-NAME or an infusion of 200 nmol/kg of S-methyl-L-cysteine for 10 minutes followed by a 500 nmol/kg bolus dose of S-methyl-L-cysteine. The measurements were then repeated a third time.

The results are summarized in Figures 1 to 6. After 72 hours of LPS exposure, blood pressure dropped 40%, heart rate dropped 15-20%, cardiac output dropped 20%, and total peripheral resistance dropped 25%. Both treatment with L-NAME and S-methyl cysteine returned mean arterial pressure to baseline values, but in the rats treated with L-NAME cardiac output, heart rate, and coronary blood flow dropped further to about 60% of the pre-treatment values while total peripheral resistance and coronary resistance increased to about 160% of baseline values. In contrast, treatment with S-methyl cysteine returned cardiac output, heart rate, total peripheral resistance, coronary blood flow, and coronary resistance to pre-treatment values. Twenty-two rats received L-NAME therapy and 7 died, while 12 rats were treated with S-methyl cysteine and none died. Twenty-eight rats received neither L-NAME nor S-methyl cysteine after treatment with lipopolysaccharide and one of those died.